

ABSTRACT OF THE DISCLOSURE

Method of identification and quantitative analysis of aldehyde(s) and/or ketone(s) in a sample by mass spectrometry using stable isotope labeled internal standard is provided. Said internal standard is prepared by reaction of an authentic sample of said aldehyde(s) and/or ketone(s) with a stable isotope labeled reagent, and is added to a sample containing said aldehyde(s) and/or ketone(s). Said aldehyde(s) and/or ketone(s) in said sample is then quantitatively converted to a chemical compound of identical structure, except the stable isotope atoms, as that of said internal standard using a non-labeled reagent. Said sample is then extracted and the extract is analyzed by mass spectrometry. Identification and quantification of said aldehyde(s) and/or ketone(s) are made from a plot of ion ratio of said converted aldehyde and/or ketone to said internal standard versus aldehyde and/or ketone concentration.

REFERENCES

US patent documents

5,559,038	September 24, 1996	J. Fred Kolhouse
6,358,996	March 19, 2002	Michael S. Alexander

Other references

Dennis J. Dietzen et al, "Facilitation of thin-layer chromatographic identification of opiates by derivatization with acetic anhydride or methoxyamine", Journal of Analytical Toxicology, Sep 1995, page 299-303, vol. 19.

Kyle R. Gee et al, "Arene chromium and manganese tricarbonyl analogs of the PCP receptor ligands 5-methyl-10,11-dihydro-5H-dibenzo[a,d]cyclohepten-5,10-imine (MK-801) and 10,5-(iminomethano)-10-11-dihydro-5H-dibenzo[a,d]cycloheptene" Journal of Organic Chemistry, 1994, p. 1492-1498, vol.59.

Hiroshi Goda et al, "Facile synthesis of 5-substituted 2-acetylthiophenes", Synthesis, 1992, p.849-851.

Arun K. Ghosh et al, "Stereoselective reduction of alpha-hydroxy oxime ethers: a convenient route to cis-1,2-amino alcohols", Tetrahedron Letters, 1991, p.711-714, vol.32.